

**EPIPEN® EPIPEN® Jr**

adrenalin 0,3/0,15mg

## Certificate for Airline Passenger (for attention of airport security and control)

### Statement for rescue medication

**EpiPen®/EpiPen® Jr** is used for the emergency treatment of sudden and life threatening allergic reactions (anaphylactic shock) to insect stings and bites, foods, drugs, latex and exercise. EpiPen®/EpiPen® Jr contains a sterile solution of adrenaline (epinephrine) for emergency intra muscular self-injection.

The patient \_\_\_\_\_ (name)

is suffering from a life threatening \_\_\_\_\_ allergy.

The EpiPen®/EpiPen® Jr (adrenaline) auto-injector is the prescribed rescue medication that the patient has to **carry at all-times**. In case of an emergency (a life threatening anaphylactic shock or a serious allergic reaction) the adrenaline can be administered and further damage caused to the patient's health can be avoided.

Physician: stamp – signature – date

### Symptoms announcing an anaphylactic reaction:

- Prickle and swelling of lips, eyes, throat and in the face
- Urticaria/itching
- Shortness of breath
- Hypotension

# EPIPEN® EPIPEN® Jr


adrenalin 0,3/0,15mg

1.



Grasp EpiPen®/EpiPen® Jr Auto-Injector in dominant hand, with thumb closest to blue safety cap. With the other hand pull off blue safety cap. **Remember: "Blue to the sky, orange to the thigh".**

2.



Hold the EpiPen®/EpiPen® Jr Auto-Injector in a distance of approximately 10 cm away from the outer thigh. The orange tip should point towards the outer thigh.

3.



Jab EpiPen®/EpiPen® Jr firmly, at a right angle (90 degree angle) to the outer thigh. Hold firmly in place for 10 seconds. The injection is now complete and the window of the Auto-Injector is obscured.

4.



The EpiPen®/EpiPen® Jr Auto-Injector should be removed and safely discarded. The orange needle cover will extend to cover the needle.

5.



Massage the injection area for 10 seconds.

## CALL EMERGENCY SERVICES IMMEDIATELY

### Application of a second EpiPen®/EpiPen® Jr:

A second dose can be given after 5-15 minutes if necessary.

### Regard the expiry date:

Replace your EpiPen®/EpiPen® Jr before reaching **the expiry date** (see «EXP») on the packaging or pen.



Meda AB

P.O. Box 906, SE-170 09 Solna.

Phone: +46 8 630 19 00. Fax: +46 8 630 19 50.

www.medasverige.se

**MEDA**